


LECTURE 10

Developing Transport Theory to Study the Chiral Phase Transition

S.P. Klevansky, A. Ogura, P. Rehberg and J. Hüfner

*Institut für Theoretische Physik, Philosophenweg 19,
D-69120 Heidelberg, Germany*



1. INTRODUCTION

One of the fundamental questions that is open to both theory and experiment is how to observe the chiral phase transition. Indications that a phase transition from a chirally symmetric to a chirally broken phase have been obtained a long time ago: lattice gauge simulations of quantum chromodynamics (QCD), which are equilibrium calculations that can include only temperature but not finite density, show a phase transition at a finite critical temperature T_c [1]. Unfortunately, due to the difficulties inherent in a confining theory such as QCD, it is impossible to observe this transition directly experimentally, let alone obtain experimental values for the usual quantities that are associated with critical phenomena, such as critical exponents.

The experimental search for clues to a phase transition that has been conceived, is to proceed via heavy ion collisions, in which both high temperatures and densities are to be reached. Much effort has been expended in this direction and will continue to be so during the early course of the next century, particularly at RHIC and CERN. However, from the theoretical side, there is at present no single conclusive observable that might indicate the occurrence of the chiral phase transition. In addition, the simple equilibrium picture of the static lattice simulations is inadequate for describing heavy ion collisions. Thus we may speculate as to whether a concrete signal of this phase transition can be obtained from a non-equilibrium formulation of the problem or transport theory, that is based on an underlying chiral Lagrangian. Here one may

surmise whether phenomena such as critical scattering, i.e. divergence of the scattering cross-sections at the phase transition, may leave an obvious signal, or simply the fact that the dynamically generated quark mass is a function of space and time may leave some visible remnant of a phase transition.

Thus our goal is to construct a consistent transport theory that is based on the Nambu–Jona-Lasinio (NJL) model, which to order $1/N_c$ also includes the mesonic degrees of freedom and the hadronization of quarks into these degrees of freedom. In this report, we discuss primarily the first term in the $1/N_c$ expansion in the collision, explicitly relating this to Feynman diagrams for quark-quark and quark-antiquark scattering (for details, see Ref.[2]), leaving the mesonic sector to a future publication [3].

2. GENERAL TRANSPORT THEORY FOR FERMIONS.

Non-equilibrium phenomena are completely described by the Schwinger - Keldysh formalism for Green functions [4]. Many conventions exist in the literature. For our purposes, it is simplest to use the convention of Landau [5]. In this, the designations $+$ and $-$ attributed to the closed time path that is shown in Fig.1, and the fermionic Green functions are defined as

$$\begin{aligned} iS^c(x, y) &= \langle T\psi(x)\bar{\psi}(y) \rangle = iS^{--}(x, y) \\ iS^a(x, y) &= \langle \tilde{T}\psi(x)\bar{\psi}(y) \rangle = iS^{++}(x, y) \\ iS^>(x, y) &= \langle \psi(x)\bar{\psi}(y) \rangle = iS^{+-}(x, y) \\ iS^<(x, y) &= -\langle \bar{\psi}(y)\psi(x) \rangle = iS^{-+}(x, y). \end{aligned} \quad (1)$$

In a standard fashion, one constructs equations of motion for the matrix of Green functions and one moves to relative and centre of mass variables, $u = x - y$ and $X = (x + y)/2$. A Fourier transform with respect to the relative coordinate, or Wigner transform,

$$S(X, p) = \int d^4u e^{ip \cdot u} S\left(X + \frac{u}{2}, X - \frac{u}{2}\right) \quad (2)$$

is then performed. Of particular interest is the equation of motion for $S^{-+} = S^{<}$. Regarding this, together with the equation of motion for the adjoint function $S^{<\dagger}$, one arrives at the so-called transport and constraint equations by adding and subtracting these. One finds

$$\frac{i\hbar}{2} \left\{ \gamma^\mu, \frac{\partial S^{-+}}{\partial X^\mu} \right\} + [\not{p}, S^{-+}(X, p)] = I_- \quad (3)$$

and

$$\frac{i\hbar}{2} \left[\gamma^\mu, \frac{\partial S^{-+}}{\partial X^\mu} \right] + \{ \not{p} - m_0, S^{-+}(X, p) \} = I_+, \quad (4)$$

respectively. Here

$$I_{\mp} = I_{\text{coll}} + I_{\mp}^A + I_{\mp}^R, \quad (5)$$

where the collision term is

$$\begin{aligned} I_{\text{coll}} &= \Sigma^{-+}(X, p) \hat{\Lambda} S^{+-}(X, p) - \Sigma^{+-}(X, p) \hat{\Lambda} S^{-+}(X, p) \\ &= I_{\text{coll}}^{\text{gain}} - I_{\text{coll}}^{\text{loss}}, \end{aligned} \quad (6)$$

and terms containing retarded (R) and advanced (A) components are contained in

$$I_{\mp}^R = -\Sigma^{-+}(X, p) \hat{\Lambda} S^R(X, p) \pm S^R(X, p) \hat{\Lambda} \Sigma^{-+}(X, p) \quad (7)$$

and

$$I_{\mp}^A = \Sigma^A(X, p) \hat{\Lambda} S^{-+}(X, p) \mp S^{-+}(X, p) \hat{\Lambda} \Sigma^A(X, p). \quad (8)$$

In these equations, Λ is given as

$$\hat{\Lambda} = \exp \left(-\frac{i\hbar}{2} \left(\frac{\overleftarrow{\partial}}{\partial X^\mu} \frac{\overrightarrow{\partial}}{\partial p_\mu} - \frac{\overleftarrow{\partial}}{\partial p_\mu} \frac{\overrightarrow{\partial}}{\partial X^\mu} \right) \right). \quad (9)$$

Note that the equations (3) and (4) are general equations that are generic for *any* relativistic fermionic theory [6].

3. APPROXIMATION SCHEMES - APPLICATION TO THE NJL MODEL

A double expansion in inverse powers of the number of colors $1/N_c$ and $\hbar$ is performed. We explain this briefly in the context of the model.

The simplest form of the NJL Lagrangian

$$\mathcal{L} = \bar{\psi} (i\hbar \not{\partial} - m_0) \psi + G \left[(\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma^5\psi)^2 \right], \quad (10)$$

is considered. Here G is a coupling strength and m_0 is the current quark mass. The color degree of freedom is implicit. As this is a strong coupling theory, an expansion in G is inadmissible, and the usual technique involves an expansion in $1/N_c$. In this ordering, the lowest order Feynman graph is a Hartree term, which describes the mean field experienced by the quarks, while higher orders introduce collisions. We examine these two cases individually. In order to make connection with the classically known transport equations, we consider an expansion of the transport and constraint equations in powers of $\hbar$.

In the transport and constraint equations, Eqs.(3) and (4), the right hand sides I_- and I_+ , respectively depend on $\hbar$ in two ways: (i) The derivative operator $\hat{\Lambda}$, Eq.(9), contains $\hbar$ to all orders. Its expansion in $\hbar$ generates the various terms of the so-called gradient expansion, and (ii) the self-energies Σ^R and Σ^{-+} may have overall factors in $\hbar$. While to order $O((1/N_c)^0)$, $\Sigma^R = m$, and is directly related to the mass, which we treat as a quantity that has a

direct classical interpretation, we find $\Sigma^{-+} \propto \hbar d\sigma/d\Omega$, and is only related via a factor $\hbar$ to a quantity with direct physical interpretation. Note that while we use the overall factors in $\hbar^0$ or $\hbar^1$ in Σ^R or Σ^{-+} in our $\hbar$ classification, we do not expand the quantities m or $d\sigma/d\Omega$ in powers of $\hbar$. We will detail this strategy in the calculation of the mean field term and collision integrals.

3.1. The Hartree Approximation

The leading term in the $1/N_c$ expansion corresponds to the Hartree approximation. The Hartree self-energy can be evaluated, given a form for the Green function. Here we assume that

$$\begin{aligned} S_H^{-+}(X, p) = & 2\pi i \frac{1}{2E_p} [\delta(p_0 - E_p) \sum_s u_s(p) \bar{u}_s(p) f_q(X, p) \\ & + \delta(p_0 + E_p) \sum_s v_s(-p) \bar{v}_s(-p) \bar{f}_{\bar{q}}(X, -p)] \end{aligned} \quad (11)$$

with $E_p^2 = p^2 + m^2(X)$ and $f_{q,\bar{q}}$ the (spin independent) quark and antiquark distribution functions. This is the quasiparticle assumption. Similar expressions can be written down for the other Green functions of Eq.(1). Inserting the quasiparticle *ansätze* into the Wigner transformed expression for the self-energy in the Hartree approximation, $-i\Sigma_H^{-+} = 2i\hbar \text{tr} i S_H^{-+}(x, x)$ leads to the non-equilibrium gap equation

$$m(X) = m_0 + 4GN_c m(X) \int \frac{d^3p}{(2\pi\hbar)^3} \frac{1}{E_p(X)} [1 - f_q(X, p) - f_{\bar{q}}(X, p)]. \quad (12)$$

Some manipulations are now necessary for the evaluation of (3) and (4). The evaluation of $I_{\pm}$ is made according to the $\hbar$ expansion. Then, on performing a spinor trace and integrating over positive or negative energies, leads one to the Vlasov equation

$$p^\mu \partial_\mu f_{q,\bar{q}}(X, \vec{p}) + m(X) \partial_\mu m(X) \partial_p^\mu f_{q,\bar{q}}(X, \vec{p}) = 0, \quad (13)$$

where $p^0 = E_p(X)$, and the constraint equation

$$(p^2 - m^2(X)) f_{q,\bar{q}} = 0. \quad (14)$$

to leading order in $\hbar$. Note that the constraint equation validates the quasiparticle *ansatz*, and that the equations Eq.(13) and (14) can be solved self-consistently.

3.2. The Collision Term

There are several graphs which contribute to the next order in the $1/N_c$ expansion. One of these is displayed in Fig. 1(a). The accompanying graph of Fig. 1(b) is of higher order in $1/N_c$, and is not required on this basis, but, as

Fig. 1. — Contributions to the self-energy that are of higher order in $1/N_c$ than the Hartree term

it turns out, is essential for a complete and correct identification of Feynman amplitudes in casting the collision term into a Boltzmann-like form. The initial task is the evaluation of I_{coll} of Eq.(6), with scalar or pseudoscalar vertices. There are three possible combinations, Σ_σ , Σ_π and Σ_{coll} , which contain unity and $i\gamma_5$ throughout or unity or $i\gamma_5$ in a mixed fashion. For example, the scalar self-energy is determined to be

$$\begin{aligned} \Sigma_\sigma^{+-}(X, p) &= -4G^2\hbar^2 \int \frac{d^4p_1}{(2\pi\hbar)^4} \frac{d^4p_2}{(2\pi\hbar)^4} \frac{d^4p_3}{(2\pi\hbar)^4} (2\pi\hbar)^4 \delta(p - p_1 + p_2 - p_3) \\ &\times [S^{+-}(X, p_1) \text{tr}(S^{-+}(X, p_2) S^{+-}(X, p_3)) \\ &\quad - S^{+-}(X, p_1) S^{-+}(X, p_2) S^{+-}(X, p_3)], \end{aligned} \quad (15)$$

while the other self-energies are similar [2].

As before, performing a spinor trace and integrating over a positive or negative energy range is required, so that the loss term, for example, is correspondingly given as

$$J_{\text{coll}, \sigma}^{q, \text{loss}} = \frac{2\pi i}{2E_p} \text{tr} \left[\Sigma_\sigma^{+-}(X, p_0 = E_p, \vec{p}) \sum_s u_s(p) \bar{u}_s(p) f_q(X, \vec{p}) \right], \quad (16)$$

which, when evaluated using the quasiparticle approximations for the Green functions, leads to eight terms. After some algebra, one can show that only terms that represent elastic scattering survive energy integration. Other terms that contain vacuum fluctuations, as well as particle production and annihilation are present at this level, see Fig. 2, but are suppressed by the quasiparticle approximation. An off-shell *ansatz* is required for them not to vanish. It is also important to note that both diagrams of Fig. 1 are essential for the identification of the Feynman amplitudes for elastic scattering in the s , t and u channels, as appropriate. While graph (a) leads to the square of the individual amplitudes, graph (b) is necessary for the mixed terms that allow one to construct

Fig. 2. — Processes of the form $q \rightarrow (q\bar{q})q$ and $(q\bar{q}) \rightarrow q$, as well as vacuum fluctuations that are suppressed in the quasiparticle approximation. These are heuristic graphs and are not Feynman graphs

a matrix element squared that involves several channels. The net result is

$$\begin{aligned}
& p^\mu \partial_\mu f_q(X, \vec{p}) + m(X) \partial_\mu m(X) \partial_p^\mu f_q(X, \vec{p}) = \\
& N_c \int d\Omega \int \frac{d^3 p_2}{(2\pi\hbar)^3 2E_{p_2}} |\vec{v}_p - \vec{v}_2| 2E_p 2E_{p_2} \\
& \times \left\{ \frac{1}{2} \frac{d\sigma}{d\Omega} \Big|_{qq \rightarrow qq} (p_2 \rightarrow 13) (f_q(p_1) \bar{f}_q(p_2) f_q(p_3) \bar{f}_q(p) - \bar{f}(p_1) f_q(p_2) \bar{f}_q(p_3) f_q(p)) \right. \\
& \left. + \frac{d\sigma}{d\Omega} \Big|_{q\bar{q} \rightarrow q\bar{q}} (p_2 \rightarrow 13) (f_q(p_1) \bar{f}_{\bar{q}}(p_2) f_{\bar{q}}(p_3) \bar{f}_q(p) - \bar{f}_q(p_1) \bar{f}_{\bar{q}}(p_3) f_{\bar{q}}(p_2) f_q(p)) \right\},
\end{aligned} \tag{17}$$

where the scattering cross-sections contain s , t and u channel graphs as appropriate, in the Born approximation and which is the relativistic generalization of the result of Kadanoff and Baym. To lowest order in $\hbar$, arguments can be given to suggest that the constraint equation remains unaltered.

4. DISCUSSION

The most important facets of this calculation are (i) in identifying precise Feynman amplitudes that contribute to the scattering processes in the collision term, and which represents the relativistic generalization of the Kadanoff-Baym

work [7], and (ii) the recognition of additional processes that would be present should a quasiparticle assumption be relaxed, such as vacuum fluctuations and particle production and annihilation. The double expansion in $\hbar$ and $1/N_c$ lays the foundation for including additional higher order terms in the $1/N_c$ expansion, from which mesons are constructed, and which enables one to build a dynamical model for both mesons and quarks. For this, we refer the interested reader to [3, 8].

We have started numerical simulations of this chiral transport theory. A first study that solves the Vlasov equation by means of a finite difference technique and which inserts collision terms in the relaxation time approximation for three flavors is presented in Ref. [9]. In this, one finds that the collisions determine the high energy parts of the particle distribution, which is exponential. The mean field, on the other hand, produces an enhancement over the exponential fits at low energies. The behavior of the quark mass as a function of time is such that a dip occurs in the initial state, and this gradually flattens as the quarks flow out.


Acknowledgments

This work has been supported in part by the Deutsche Forschungsgemeinschaft DFG under the contract number Hu 233/4-4, and by the German Ministry for Education and Technology under contract number 06 HD 742.

References


- [1] See for example, Laermann E., *Nucl. Phys.* **A610** (1996) 1.
- [2] Klevansky S.P., Ogura A., and Hüfner J., Heidelberg Preprint HD-TVP-9702.
- [3] Klevansky S.P., Rehberg P., and Hüfner J., in preparation.
- [4] Schwinger J., *J. Math. Phys.* **2** (1961) 407. Keldysh L.V., *JETP* **20** (1965) 1018.
- [5] Lifschitz E.M. and Pitajewski L.P., *Physikalische Kinetik* (Akademie Verlag, Berlin, 1986) p. 423.
- [6] See for example, Mrówczyński S. and Heinz U., *Ann. Phys. (N.Y.)* **229** (1994) 1.
- [7] Kadanoff L.P. and Baym G., *Quantum Statistical Mechanics* (Benjamin / Cummings, Massachusetts, 1962) p. 110.
- [8] Klevansky S.P., Rehberg P., Ogura A., and Hüfner J., *QCD Phase Transitions* (GSI, Darmstadt, 1997) p. 397.
- [9] Rehberg P. and Hüfner J., Heidelberg Preprint HD-TVP-97/03

$$-i\Sigma^{+-} =$$



(a)

$+$



(b)

